

Week 3 Abstracts: Sept. 8-12, 2025

Speaker: Ruobing Zhang

Abstract: A Poincaré-Einstein manifold is a complete non-compact Einstein manifold with negative scalar curvature which can be conformally deformed to a compact manifold with boundary, called the conformal boundary or conformal infinity. Naturally, such a space is associated with a conformal structure on the conformal infinity. A fundamental theme in studying these geometric objects is to relate the Riemannian geometric data of the Einstein metric to the conformal geometric data at infinity which is also called the AdS/CFT correspondence in theoretical physics. In this talk, we will explore some new techniques from the metric geometric point of view, by which one can establish some new rigidity, quantitative rigidity, and regularity results.

Speaker: Ariadna Leon Quiros

Abstract: In 1977, D. C. Robinson developed a method for proving static vacuum Black Hole uniqueness in General Relativity. This method has recently been generalized to higher dimensions by C. Cederbaum, A. Cogo, B. Leandro, and J. Paolo dos Santos. It turns out that the same philosophy can prove several geometric inequalities. In my talk, I will show how to adjust this approach to prove the Riemannian Positive Mass Theorem. If time permits, I will discuss other related results. This is joint work with C. Cederbaum and B. Meco.

Speaker: Roland Steinbauer

Abstract: The cut-and-paste method is a procedure for constructing null thin shells by matching two regions of the same spacetime across a null hypersurface. Originally proposed by Penrose, it has so far allowed to describe purely gravitational and null-dust shells in constant-curvature backgrounds. In this paper, we extend the cut-and-paste method to null shells with arbitrary gravitational/matter content. To that aim, we first derive a locally Lipschitz continuous form of the metric of the spacetime resulting from the most general matching of two constant-curvature spacetimes with totally geodesic null boundaries, and then obtain the coordinate transformation that turns this metric into the cut-and-paste form with a Dirac-delta term. The paper includes an example of a null shell with non-trivial energy density, energy flux and pressure in Minkowski space. <https://arxiv.org/abs/2508.00231> This is joint work with Miguel Manzano and Argam Ohanyan.

Speaker: Andrea Mondino

Abstract: TBA

Speaker: Guoliang Yu

Abstract: In this talk, I will introduce the higher index theory of elliptic differential operators and discuss recent developments and applications in geometry and topology. I will make an effort to keep the presentation accessible to graduate students.

Speaker: Zhizhang Xie

Abstract: In this talk, I will present my joint work with Jinmin Wang and Guoliang Yu on a new index theorem for manifolds with singularities, such as manifolds with corners and, more generally, manifolds with polyhedral-type boundary. As an application, we obtained a positive solution to Gromov's dihedral rigidity conjecture. This conjecture concerns comparisons of scalar curvature, mean curvature and dihedral angles for compact manifolds with polyhedral-type boundary, and has very interesting implications in geometry and mathematical physics. Further developments of this new index theorem have led us to a positive solution of Gromov's flat corner domination conjecture. As a consequence, we answered positively a long standing conjecture in discrete geometry — the Stoker conjecture.

Speaker: Gilbert Weinstein

Abstract: In this talk, I will discuss our recent result. Either there is a counterexample to black hole uniqueness, or the following statement holds. Axisymmetric, complete, simply connected, maximal initial data sets for the Einstein equations of nonnegative energy density with ends that are either asymptotically flat or asymptotically cylindrical, admit an ADM mass lower bound given by the square root of total angular momentum. Moreover, equality is achieved only for a constant time slice of an extreme Kerr spacetime. The proof is based on a novel flow of singular harmonic maps with hyperbolic plane target, under which the renormalized harmonic energy is monotonically nonincreasing. Relevant properties of the flow are achieved through a refined asymptotic analysis of solutions to the linearized harmonic map equations. This is joint work with Qing Han, Marcus Khuri, and Jingang Xiong.

Speaker: Roya Mohayaee

Abstract: Monge-Ampère equation, a nonlinear generalization of the Poisson equation, provides an alternative nonrelativistic description of gravity. The equation is affine invariant, has rich geometric properties, connects to optimal transport theory, and remains bounded at short distances. Monge-Ampère gravity naturally emerges through the application of large-deviation principle to a Brownian system of indistinguishable and independent particles. I show that Monge-Ampère gravity can describe a scalar field, often evoked in modified theories of gravity such as Galileons. We show that Monge-Ampère gravity, as a nonlinear model of a new scalar field, is screened at short distances, and behaves differently from Newtonian gravity above certain scales but approaches it asymptotically. I show how the Monge-Ampère equation can be obtained from the fully covariant Lagrangian of Galileon theories in the static limit. I finally present the numerical results from N-body simulations of this model based on semi-discrete optimal transport algorithm.